


# Equilibria

## A-Level Chemistry

### Conjugate acids and bases

When a Brønsted acid loses an  $H^+$ , what is left is its **conjugate base** 共轭碱. When a base gains an  $H^+$ , it becomes its **conjugate acid** 共轭酸. The two species that differ by just one  $H^+$  form a **conjugate acid–base pair** 共轭酸碱对.

For example, in  $CH_3COOH \rightleftharpoons CH_3COO^- + H^+$ , the pair is  $CH_3COOH$  (acid) and  $CH_3COO^-$  (its conjugate base).




*Conjugate pairs differ by one proton: an acid loses  $H^+$  to give its conjugate base; a base gains  $H^+$  to give its conjugate acid*

## pH and the equilibrium constants

- **pH** measures acidity:  $\text{pH} = -\log[\text{H}^+]$ , so  $[\text{H}^+] = 10^{-\text{pH}}$ .
- the **acid dissociation constant** 酸解离常数 of a weak acid HA is  $K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$ , and  $\text{p}K_a = -\log K_a$ . A larger  $K_a$  (smaller  $\text{p}K_a$ ) means a stronger acid.
- the **ionic product of water** 水的离子积 is  $K_w = [\text{H}^+][\text{OH}^-] = 1.0 \times 10^{-14}$  at 298 K.

$$\text{pH} = -\log[\text{H}^+]$$



*The pH scale:  $\text{pH} = -\log$  of the hydrogen-ion concentration*



*pH paper estimates the pH of a solution from the colour it turns*

## Calculating pH

- **strong acid** 强酸: fully ionised, so  $[\text{H}^+]$  equals the acid concentration; then take  $-\log$ .
- **strong alkali** 强碱: find  $[\text{OH}^-]$  from the concentration, then use  $[\text{H}^+] = K_w/[\text{OH}^-]$ .
- **weak acid** 弱酸: only partly ionised, so use  $[\text{H}^+] = \sqrt{K_a \times [\text{HA}]}$ .

**Worked example.** Find the pH of  $0.050 \text{ mol dm}^{-3}$  hydrochloric acid (a strong acid).

It is fully ionised, so  $[\text{H}^+] = 0.050 \text{ mol dm}^{-3}$ , giving  $\text{pH} = -\log(0.050) = 1.30$ .

**Worked example.** Find the pH of  $0.10 \text{ mol dm}^{-3}$  sodium hydroxide (a strong base). ( $K_w = 1.0 \times 10^{-14}$ .)

$[\text{OH}^-] = 0.10$ , so  $[\text{H}^+] = K_w/[\text{OH}^-] = (1.0 \times 10^{-14})/0.10 = 1.0 \times 10^{-13}$ , giving  $\text{pH} = 13.0$ .

**Worked example.** Find the pH of  $0.10 \text{ mol dm}^{-3}$  ethanoic acid (a weak acid). ( $K_a = 1.8 \times 10^{-5}$ .)


$$[\text{H}^+] = \sqrt{K_a \times [\text{HA}]} = \sqrt{(1.8 \times 10^{-5})(0.10)} = 1.3 \times 10^{-3}, \quad \text{pH} = -\log(1.3 \times 10^{-3}) = 2.87.$$

# Buffer solutions

A **buffer solution** 缓冲溶液 resists a change in pH when a small amount of acid or alkali is added. You make one from a weak acid and its conjugate base (for example ethanoic acid and sodium ethanoate).

It works because the mixture holds a store of both partners:

- added  $\text{H}^+$  is removed by the conjugate base:  $\text{CH}_3\text{COO}^- + \text{H}^+ \rightarrow \text{CH}_3\text{COOH}$ .
- added  $\text{OH}^-$  is removed by the weak acid:  $\text{CH}_3\text{COOH} + \text{OH}^- \rightarrow \text{CH}_3\text{COO}^- + \text{H}_2\text{O}$ .



either way, the pH barely changes

*A buffer holds a store of a weak acid and its conjugate base: added  $\text{H}^+$  is mopped up by  $\text{A}^-$  and added  $\text{OH}^-$  by  $\text{HA}$ , so the pH barely changes*

To find the pH, put the concentrations of the acid and its salt into the  $K_a$  expression. Buffers are important in living things — for example,  $\text{HCO}_3^-$  keeps the pH of blood close to 7.4.

## Solubility product

For a salt that barely dissolves, the **solubility product** 溶度积 ( $K_{\text{sp}}$ ) is the product of the ion concentrations in a saturated solution, each raised to the power of its number in the formula:

$$K_{\text{sp}} = [\text{Ag}^+][\text{Cl}^-] \quad K_{\text{sp}} = [\text{Ca}^{2+}][\text{F}^-]^2$$

You can find  $K_{\text{sp}}$  from the solubility, or the solubility from  $K_{\text{sp}}$ .



*Stalactites grow as dissolved calcium carbonate slowly comes back out of solution — a real solubility equilibrium*

## The common ion effect


The **common ion effect** 同离子效应 is the way a salt becomes **less** soluble in a solution that already contains one of its ions. The extra ion pushes the dissolving equilibrium back (Le Chatelier), so less salt dissolves. You can calculate the new solubility using  $K_{\text{sp}}$  and the concentration of the common ion.

## Partition coefficients

When a solute is shaken with two solvents that do not mix, it spreads between them. The **partition coefficient** 分配系数 ( $K_{\text{pc}}$ ) is the ratio of its concentrations in the two layers (at constant temperature):

$$K_{\text{pc}} = \frac{[\text{solute in solvent 1}]}{[\text{solute in solvent 2}]}$$

This works when the **solute** 溶质 is in the same physical state in both solvents. The value depends on the **polarity** 极性 of the solute and of each **solvent** 溶剂: a non-polar solute dissolves more in the non-polar solvent, while a polar solute prefers the polar solvent.



*A solute shaken with two immiscible solvents spreads between them; the partition coefficient is the ratio of its concentrations in the two layers*

## Exam tips

- $\text{pH} = -\log[\text{H}^+]$ ; for a **strong acid**  $[\text{H}^+] = \text{concentration}$ , for a **weak acid** use  $[\text{H}^+] = \sqrt{K_a[\text{HA}]}$ .
- For a base, get  $[\text{H}^+]$  from  $K_w = [\text{H}^+][\text{OH}^-] = 1.0 \times 10^{-14}$ .
- For a **buffer** use  $[\text{H}^+] = K_a \times [\text{acid}]/[\text{salt}]$  and explain how it removes added  $\text{H}^+/\text{OH}^-$ .
- Write the  $K_{sp}$  expression with the correct powers; the number of decimal places in a pH equals the significant figures of  $[\text{H}^+]$ .