

Exercise walk-through

Polar coordinates ■ 1.5

eXercise

You must be able to


- convert between (r, θ) and (x, y)
- sketch a simple polar curve (e.g. a cardioid)
- find an area with $\frac{1}{2} \int r^2 d\theta$

Method — Conversion & area

Convert $r = 4 \cos \theta$: $\times r \rightarrow r^2 = 4r \cos \theta$, so $x^2 + y^2 = 4x$, i.e. $(x - 2)^2 + y^2 = 4$ (circle).


Area: $A = \frac{1}{2} \int_{\alpha}^{\beta} r^2 d\theta$.

Method — Conversion & area



A point reached by distance r at angle θ , with x and y components

Method — Conversion & area



cardioid · polar area $A = \frac{1}{2} \int_{\alpha}^{\beta} r^2 d\theta$

Polar curves are traced as the angle varies

Practice 2.1 ■ 2 marks

Convert the point $(r, \theta) = (2, \frac{\pi}{3})$ to Cartesian coordinates.

Solution 2.1

Method: Conversion & area

Worked steps

$$x = 2 \cos \frac{\pi}{3} = 1, y = 2 \sin \frac{\pi}{3} = \sqrt{3} \quad \text{M1} \quad \text{A1}.$$

Practice 2.2 ■ 1 mark

Write down the formula for the area of a polar sector.

Solution 2.2

Method: Conversion & area

Worked steps

$$A = \frac{1}{2} \int r^2 d\theta \quad \text{B1}.$$

Exam-style 3.1 ■ 1 mark

The polar equation $r = 2 \sin \theta$ represents:

- A** a line
- B** a circle
- C** a parabola
- D** a spiral

Solution 3.1

Method: Conversion & area

A a line

B a circle



C a parabola

D a spiral

Worked steps

B. $r = 2 \sin \theta \Rightarrow r^2 = 2r \sin \theta \Rightarrow x^2 + y^2 = 2y$, a circle.

Exam-style 3.2 ■ 2 marks

A curve has polar equation $r = 1 + \cos \theta$ (a cardioid).

(a) Find r when $\theta = 0$ and $\theta = \frac{\pi}{2}$.

(b) Find the area enclosed by the curve, using $A = \frac{1}{2} \int_0^{2\pi} (1 + \cos \theta)^2 d\theta$.

Solution 3.2

Method: Conversion & area

Worked steps

(a) $\theta = 0: r = 2; \theta = \frac{\pi}{2}: r = 1$ **B1** **B1**.

(b) $(1 + \cos \theta)^2 = 1 + 2 \cos \theta + \cos^2 \theta = \frac{3}{2} + 2 \cos \theta + \frac{1}{2} \cos 2\theta$ **M1**; $A = \frac{1}{2} \left[\frac{3}{2} \theta + 2 \sin \theta + \frac{1}{4} \sin 2\theta \right]_0^{2\pi}$ **M1** **M1**; $= \frac{1}{2} \left(\frac{3}{2} \cdot 2\pi \right) = \frac{3\pi}{2}$ **M1** **A1**.

Exam-style 3.3 ■ 4 marks

Convert $r = \frac{2}{1 + \cos \theta}$ to Cartesian form.

Solution 3.3

Method: Conversion & area

Worked steps

$$r + r \cos \theta = 2 \Rightarrow r = 2 - x \text{ (since } r \cos \theta = x \text{) } \text{M1}; r^2 = (2 - x)^2 \Rightarrow x^2 + y^2 = 4 - 4x + x^2 \text{ M1 M1}; y^2 = 4 - 4x, \text{ i.e. } y^2 = -4(x - 1) \text{ A1}.$$