

3.5 Linear motion under a variable force

Name: _____ Class: _____ Date: _____

Total: 14 marks

KEY WORDS differential equation 微分方程 • variable force 变力

Objective


Build the skills to answer exam questions on **linear motion under a variable force** 变力下的直线运动 — using $a = v \frac{dv}{dx}$ (force depends on position) or $a = \frac{dv}{dt}$ (force depends on time). Take $g = 10 \text{ m s}^{-2}$.

You must be able to:

- choose $a = v \frac{dv}{dx}$ for a position-dependent force
- set up and solve the resulting differential equation
- use $a = \frac{dv}{dt}$ for a time-dependent force

1 Worked examples

■ Force depending on position



A position-dependent force (e.g. a spring) makes Newton's law a differential equation in v and x

8 kg under force $(x^3 + 4x)$ N, $v = 1$ at $x = 0$: $8v \frac{dv}{dx} = x^3 + 4x$; integrate $\rightarrow 4v^2 = \frac{1}{4}x^4 + 2x^2 + 4$, so $v = \frac{1}{4}x^2 + 1$.

2 Practice

2.1 State the form of acceleration to use when a force depends on **position** x . [1]

2.2 A particle has $v \frac{dv}{dx} = 2x$. Find v^2 in terms of x , given $v = 0$ at $x = 0$. [2]

3 Exam-style questions

3.1 For a force that depends on position, acceleration is best written as: [1]

- A $\frac{dv}{dt}$
 - B $v \frac{dv}{dx}$
 - C $\frac{dx}{dt}$
 - D $\frac{d^2x}{dt^2}$ only
-

3.2 A particle of mass 2 kg moves along a straight line under a force $F = 6x$ N in the direction of motion. At $x = 0$ the speed is 2 m s^{-1} .

(a) Form a differential equation in v and x . [2]

(b) Find v when $x = 2$. [4]

3.3 A particle of mass 0.5 kg moves so that its acceleration is $\frac{dv}{dt} = 4 - 2t$. At $t = 0$ it is

at rest. Find its velocity when $t = 3$ and comment.

[4]

Solutions

2.1 $a = v \frac{dv}{dx}$.

2.2 $\int v \, dv = \int 2x \, dx \Rightarrow \frac{1}{2}v^2 = x^2 + C$; $v = 0$ at $x = 0 \Rightarrow C = 0$, so $v^2 = 2x^2$.

3.1 B. Use $a = v \frac{dv}{dx}$ for a position-dependent force.

3.2 (a) $2v \frac{dv}{dx} = 6x$, i.e. $v \frac{dv}{dx} = 3x$.

(b) $\int v \, dv = \int 3x \, dx \Rightarrow \frac{1}{2}v^2 = \frac{3}{2}x^2 + C$; at $x = 0$, $v = 2 \Rightarrow C = 2$; at $x = 2$: $\frac{1}{2}v^2 = 6 + 2 = 8 \Rightarrow v^2 = 16$; $v = 4 \text{ m s}^{-1}$.

3.3 $v = \int (4 - 2t) \, dt = 4t - t^2 + C$; $v = 0$ at $t = 0 \Rightarrow C = 0$; at $t = 3$: $v = 12 - 9 = 3 \text{ m s}^{-1}$; the force has reversed (since $4 - 2t < 0$ for $t > 2$), so the particle is slowing after $t = 2$.