

A-Level Physics

Name: _____ Score: _____

1. The rate of change of velocity with time is called _____.

2. A car starts at $u = 2 \frac{\text{m}}{\text{s}}$ and accelerates at $a = 3 \frac{\text{m}}{\text{s}^2}$ for $t = 4 \text{ s}$. Find v .
_____ m/s
3. A ball thrown horizontally and a ball dropped from the same height hit the ground at the same time (no air resistance).
☐ True ☐ False
4. Which quantity is the **rate of change of displacement**?
☐ speed
☐ velocity
☐ acceleration
☐ distance
5. Which equation is just the **gradient** of a velocity–time graph?
☐ $v = u + at$
☐ $s = ut + \frac{1}{2}at^2$
☐ $s = \frac{1}{2}(u + v)t$
☐ $v^2 = u^2 + 2as$
6. A projectile is launched at $20 \frac{\text{m}}{\text{s}}$, 30° above the horizontal. What is the **vertical** part of the launch velocity?
_____ m/s
7. Deceleration is a completely separate quantity from acceleration.
☐ True ☐ False
8. With no air resistance, a heavy ball falls faster than a light one.
☐ True ☐ False
9. Select **all** the statements that are true for a projectile (ignore air).
☐ The horizontal velocity stays constant
☐ The vertical velocity is zero at the highest point

- ☐ The whole velocity is zero at the highest point
- ☐ The path is a parabola
- ☐ A heavier projectile falls faster

Tick every correct option.

10. On a displacement–time graph, a flat (horizontal) line means the object is at rest.

- ☐ True ☐ False

A-Level Physics

1. **acceleration**

Acceleration is how fast the velocity changes — a vector, in $\frac{\text{m}}{\text{s}^2}$.

2. **14 m/s**

Use $v = u + at = 2 + 3 \times 4 = 14 \frac{\text{m}}{\text{s}}$.

3. **True**

Yes — the vertical motion is the same free fall for both, and the horizontal speed does not change the fall time.

4. **velocity**

Velocity is the rate of change of *displacement* (a vector). Speed is the rate of change of *distance* (a scalar).

5. $v = u + at$

The gradient is $a = \frac{v - u}{t}$, which rearranges to $v = u + at$.

6. **10 m/s**

Vertical part $= u \sin \theta = 20 \times \sin 30^\circ = 20 \times 0.5 = 10 \frac{\text{m}}{\text{s}}$.

7. **False**

No — deceleration is just acceleration in the opposite direction to the motion (a negative acceleration).

8. **False**

No — free fall has the same acceleration g for every mass. Air resistance is what makes a feather seem to fall slower.

9. **The horizontal velocity stays constant; The vertical velocity is zero at the highest point; The path is a parabola**

At the top only the *vertical* velocity is zero — the horizontal velocity $u \cos \theta$ carries on, so the ball keeps moving. Mass does not change free fall.

10. **True**

A flat line has zero gradient, and the gradient is the velocity — so the velocity is zero (at rest).