


Ideal Gases

A2 Physics ■ Topic 15

A-Level Physics



Ideal Gases

What we will cover

- 1 The mole & N_A
- 2 $pV = nRT = NkT$
- 3 The gas laws
- 4 Kinetic theory
- 5 Molecular KE = $\frac{3}{2}kT$
- 6 Recap



1 The mole

Ideal Gases

└ The mole

The mole

Amount of substance 物质的量 n is measured in **moles** 摩尔. One mole holds **Avogadro constant** 阿伏伽德罗常数 particles – **atoms** 原子 for a **monatomic** 单原子 gas like helium, molecules for O_2 :

$$N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$$


Number of particles $N = nN_A$. Molar mass in grams is the relative formula mass – 18 g of water is one mole.

Count particles with moles, not with 10^{23} on every line. n is the quantity the gas laws actually use.

Ideal Gases

└ The mole

One mole, counted



$$= 6.02 \times 10^{23} \text{ particles } (N_A)$$

1 mole

A mole is 6.02×10^{23} particles – the number chosen so that a mole of a substance weighs its relative molecular mass in grams.

One mole is 6.02×10^{23} particles – whatever the particle is. The molar mass in grams equals the relative formula mass, so 18 g of water is exactly one mole.

2 Equation of state

Ideal Gases
└ Equation of state

The equation of state


An **ideal gas** 理想气体 obeys $pV \propto T$ exactly:
$$pV = nRT \quad \text{or} \quad pV = NkT$$

- the **molar gas constant** 摩尔气体常数 $R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$ – per **mole**
- the **Boltzmann constant** 玻尔兹曼常数 $k = 1.38 \times 10^{-23} \text{ J K}^{-1}$ – per **molecule**
- $N = nN_A$, so $k = R/N_A$

Match the form to your **amount**: moles $\rightarrow nRT$, molecules $\rightarrow NkT$. SI through-out – Pa, m^3 , and **kelvin**.

Ideal Gases
└ Equation of state

The equation of state, in detail


$$pV = nRT$$

gas: p , V , T , n moles

Every quantity in $pV = nRT$ is in SI units: pressure in Pa, volume in m^3 , and temperature in **kelvin** – never $^{\circ}\text{C}$.

Ideal Gases
└ Equation of state

Worked example – how many moles?

A 0.020 m^3 cylinder holds gas at 27°C and $2.0 \times 10^5 \text{ Pa}$. Find n . ($R = 8.31$)

First to kelvin: $T = 27 + 273 = 300 \text{ K}$.
$$n = \frac{pV}{RT} = \frac{(2.0 \times 10^5)(0.020)}{8.31 \times 300} \approx 1.6 \text{ mol}$$

The kelvin conversion is the **first** line, every time. 27 in place of 300 would inflate n by a factor of 11.

3

The gas laws

Ideal Gases
└ The gas laws

A fixed mass changing state

For a fixed amount of gas:
$$\frac{p_1 V_1}{T_1} = \frac{p_2 V_2}{T_2}$$
Hold one quantity fixed and a named law drops out.

Boyle
 T fixed: pV constant

Charles
 p fixed: V/T constant


pressure law
 V fixed: p/T constant

$pV \propto T$ holds **only** in kelvin – $^{\circ}\text{C}$ here is the classic lost mark.

Ideal Gases
└ The gas laws

Boyle's law

At constant temperature pV is constant, so p against $1/V$ is a **straight line through the origin** – the usual way an exam asks you to verify it.




The graph shows pressure p on the vertical axis and volume V on the horizontal axis. Two hyperbolic curves represent isotherms. The lower curve is labeled T_1 and the upper curve is labeled T_2 with the text 'higher T_2 '. A dashed line segment is tangent to the T_2 curve, with the equation $pV = \text{const (T fixed)} \Rightarrow p \propto 1/V$ written next to it.

Charles's law and absolute zero

At constant pressure the volume climbs **linearly** with temperature. Extrapolate the line back to **zero volume** and it meets $\approx -273^\circ\text{C}$.

Every gas points at the **same** intercept – which is what makes absolute zero a property of **temperature**, not of any one gas.



$V \propto T$ · Charles; extrapolates to absolute zero

Worked example – heating at constant pressure

A fixed mass at 300 K fills 0.50 m^3 . Heat it to 450 K at constant pressure. Find the new volume.

p is fixed, so V/T is constant:

$$V_2 = V_1 \frac{T_2}{T_1} = 0.50 \times \frac{450}{300} = 0.75\text{ m}^3$$

Cancel what is **constant** before substituting – here p drops out and only the temperature **ratio** matters.


4 Kinetic theory

Kinetic theory: the assumptions

- 1 many identical molecules in **random motion** 无规则运动
- 2 their own volume is negligible
- 3 collisions are brief; straight lines between
- 4 no **intermolecular** 分子间 forces except in a collision
- 5 collisions are **elastic** 弹性碰撞
- 6 Newton's laws apply

Elastic is what keeps a gas from **cooling** itself. The model fails at **high** p (volume matters) and **low** T (forces matter).

Assumptions




1. many identical molecules, in continuous **random motion**
2. their own volume is **tiny** next to the container
3. straight lines between hits - intermolecular forces **ignored**
4. collisions are **elastic**: no kinetic energy is lost

poor at very high pressure (volume matters) or very low temperature (forces matter)

The key assumptions of the kinetic theory of an ideal gas

Where the pressure comes from



One molecule, one wall, in a cube of side L :

- bounce: the **momentum** 动量 change is $\Delta p_x = -2mu_{x1}$, so the wall takes an **impulse** 冲量 of $+2mu_{x1}$
- next hit after $\Delta t = 2L/u_{x1}$
- so $F_1 = \frac{\Delta p}{\Delta t} = \frac{mu_{x1}^2}{L}$

Sum over N , then use $\langle u_x^2 \rangle = \frac{1}{3} \langle c^2 \rangle$:

$$pV = \frac{1}{3} Nm \langle c^2 \rangle$$


The $\frac{1}{3}$ is **symmetry**: no direction is special, so each axis carries a third of the motion.

Root-mean-square speed

r.m.s. speed 均方根 $= \sqrt{\langle c^2 \rangle}$: square each speed, average, then square-root. It sits slightly **above** the mean speed – squaring weights the fast molecules more.

The gas needs $\langle c^2 \rangle$, not $\langle c \rangle$, because **energy** goes as c^2 . Averaging the speeds first gives the wrong answer.

Root-mean-square speed, in detail



read the name **backwards**: root of the mean of the squares - squaring weights fast molecules more, so c_{rms} is slightly above the mean speed

Square every speed, average, then square-root – in that order. Averaging the speeds first gives a different, smaller number, because the fast molecules count for more once squared.

Why it matters under water




A diver's air is compressed: at 10 m the pressure is doubled, so a given mass of gas takes half the volume.

The spread of speeds

Molecules do not share one speed – they occupy a **distribution**. Raise the temperature and the curve **broadens** and shifts to faster speeds.

The area is fixed (the molecules still all exist), so a broader curve must also be a **lower** one.

The spread of speeds, in detail



The spread of speeds Molecules do not share one speed – they occupy a distribution.

5 Molecular kinetic energy

Average translational KE

The **average translational kinetic energy** 平均平动动能, from two expressions for the same pV – the second using the **mean square** 均方 speed $\langle c^2 \rangle$:


$$pV = NkT, \quad pV = \frac{1}{3}Nm\langle c^2 \rangle$$

Set equal, cancel N , multiply by $\frac{3}{2}$:

$$\langle E_k \rangle = \frac{1}{2}m\langle c^2 \rangle = \frac{3}{2}kT$$

The average molecular KE depends on **temperature alone** – not the gas, not the pressure. Squeezing a gas at fixed T gives more wall hits, **not** faster molecules.

Average translational KE, in detail



Mean molecular kinetic energy depends on **T alone**. Squeezing a gas at fixed T gives more wall hits, not faster molecules.

Consequences

- **doubling the absolute temperature doubles the average KE** of each molecule, so $\langle c^2 \rangle$ doubles and $c_{r.m.s.}$ grows by $\sqrt{2}$.
- for **two gases at the same temperature**, the lighter gas has a larger $\langle c^2 \rangle$. Hydrogen molecules move faster on average than oxygen molecules in the same room.
- **total translational KE** of N molecules: $\frac{3}{2}NkT = \frac{3}{2}nRT$.

Worked example – r.m.s. speed of oxygen

Find $c_{r.m.s.}$ for O_2 at 300 K.

$$m = 5.3 \times 10^{-26} \text{ kg}, \quad k = 1.38 \times 10^{-23} \text{ J K}^{-1}$$

$$\text{From } \frac{1}{2}m\langle c^2 \rangle = \frac{3}{2}kT: \quad \langle c^2 \rangle = 3kT/m$$

$$c_{r.m.s.} = \sqrt{\frac{3(1.38 \times 10^{-23})(300)}{5.3 \times 10^{-26}}}$$

$$\approx 480 \text{ m s}^{-1}$$


Faster than a jet – yet the gas goes nowhere, because the **directions** are random. Lighter gas, same $T \Rightarrow$ **faster**: hydrogen outruns oxygen in the same room.

Internal energy of an ideal gas

Ideal molecules are points: no intermolecular PE, no rotation or vibration in this model. So the **internal energy** 内能 is **all** kinetic:

$$U = \frac{3}{2}NkT = \frac{3}{2}nRT$$

$U \propto T$ – double the thermodynamic temperature and you double the internal energy.



Exam tips

- Use $pV = nRT$ (n in mol) or $pV = NkT$ (N molecules), with temperature in **kelvin**.
- Learn the **kinetic-theory assumptions** (random motion, negligible molecular volume, elastic collisions, no intermolecular forces).
- Mean translational KE = $\frac{3}{2}kT$ – it depends **only on temperature**.

6

Recap

Ideal Gases
↳ Recap

Topic 15 – key takeaways

1 The mole

$$N = nN_A, N_A = 6.02 \times 10^{23}$$

2 State

$$pV = nRT = NkT, \text{ always in kelvin}$$

3 Kinetic theory

$$pV = \frac{1}{3}Nm\langle c^2 \rangle \text{ from elastic bounces}$$


4 Energy

$$\langle E_k \rangle = \frac{3}{2}kT - \text{temperature only}$$

Ideal Gases
↳ Recap

Check yourself

- 1 How many molecules in 2.0 mol?
- 2 Which form suits a count of molecules?
- 3 Why is there a $\frac{1}{3}$ in $pV = \frac{1}{3}Nm\langle c^2 \rangle$?
- 4 At the same T , which is faster: H_2 or O_2 ?



Answers 1. 1.2×10^{24} 2. $pV = NkT$ 3. symmetry – a third per axis 4. H_2