

1.13 Removing Discontinuities

Name: _____ Class: _____ Date: _____

Total: 9 marks

KEY WORDS removable discontinuity 可去间断点 • hole 空洞 • limit 极限 • continuous 连续 • jump 跳跃

Objective

Build the skills to answer exam questions on **removing discontinuities**.**You must be able to:**

- remove a **removable discontinuity** 可去间断点 by redefining f at the hole

1 Worked examples

Study these first. Each one shows the method for a question type used later.

■ Removing a discontinuity

A removable discontinuity (a hole) can be removed by defining $f(a) = \lim_{x \rightarrow a} f(x)$.

■ Example

 $f(x) = \frac{x^2 - 4}{x - 2}$ has a hole at $x = 2$; since the limit there is 4, setting $f(2) = 4$ makes f continuous.

2 Practice

2.1 State how to remove a removable discontinuity. [1]

2.2 For $f(x) = \frac{x^2 - 4}{x - 2}$, find the value that removes the hole at 2. [2]

2.3 State which kind of discontinuity can be removed. [1]

3 Exam-style questions

3.1 A removable discontinuity is removed by setting $f(a)$ equal to [1]

- **A** 0
 - **B** the limit at a
 - **C** $f(a + 1)$
 - **D** ∞
-

3.2 Which discontinuity can be removed? [1]

- **A** jump
 - **B** infinite
 - **C** removable
 - **D** none
-

3.3 $f(x) = \frac{x^2 - 9}{x - 3}$, with a hole at 3.

(a) Simplify f . [1]

(b) Find the limit at 3. [1]

(c) State $f(3)$ to remove the hole. [1]

Solutions

2.1 define $f(a)$ to equal the limit $\lim_{x \rightarrow a} f(x)$.

2.2 the limit is 4, so set $f(2) = 4$.

2.3 a removable discontinuity (a hole).

3.1 B.

3.2 C.

3.3 (a) $x + 3$ (for $x \neq 3$). (b) 6. (c) $f(3) = 6$.