

10.7 Alternating Series Test for Convergence

Name: _____ Class: _____ Date: _____

Total: 11 marks

KEY WORDS alternating series 交错级数 • converges 收敛 • diverges 发散 • harmonic series 调和级数
• partial sums 部分和

Objective

Build the skills to answer exam questions on the **alternating series test**.

You must be able to:

- recognise an **alternating series** 交错级数 $\sum(-1)^n b_n$
- check the three conditions ($b_n > 0$, decreasing, $\rightarrow 0$)
- conclude convergence when they hold

1 Worked examples

Study these first. Each one shows the method for a question type used later — follow the steps and you can do the Practice and Exam-style questions yourself.

■ The alternating series test

An alternating series $\sum(-1)^n b_n$ (with $b_n > 0$) **converges** if:

1. b_n is **decreasing** ($b_{n+1} \leq b_n$), and
2. $\lim_{n \rightarrow \infty} b_n = 0$.

■ A worked example

$\sum \frac{(-1)^n}{n}$: $b_n = \frac{1}{n}$ is positive, decreasing, and $\rightarrow 0$. So the **alternating harmonic series converges** — even though $\sum \frac{1}{n}$ diverges.

■ Both conditions matter

If the terms b_n do **not** go to 0, the n th term test already forces divergence. If $b_n \rightarrow 0$ but is not decreasing, the test does not directly apply.

■ Recognising the form

The sign alternates via $(-1)^n$ or $(-1)^{n+1}$. Peel off the sign to get b_n , then test b_n .

2 Practice

Now apply the methods above.

2.1 For $\sum \frac{(-1)^n}{n^2}$, identify b_n . [1]

2.2 Check the two conditions for $\sum \frac{(-1)^n}{n^2}$. [2]

2.3 State whether $\sum \frac{(-1)^n}{n^2}$ converges. [1]

3 Exam-style questions

3.1 The alternating series test requires that b_n is positive, tends to 0, and is [1]

- **A** increasing
 - **B** decreasing
 - **C** constant
 - **D** geometric
-

3.2 Consider $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{\sqrt{n}}$.

(a) Identify b_n and check the conditions. [3]

(b) State the conclusion. [1]

3.3 Explain why the alternating harmonic series $\sum \frac{(-1)^n}{n}$ converges even though the

harmonic series diverges.

[2]

Solutions

2.1 $b_n = \frac{1}{n^2}$.

2.2 $b_n = \frac{1}{n^2} > 0$ and decreasing; $\lim b_n = 0$.

2.3 Converges.

3.1 B — b_n must be decreasing.

3.2 (a) $b_n = \frac{1}{\sqrt{n}} > 0$, decreasing, and $\rightarrow 0$. (b) Converges by the alternating series test.

3.3 The alternating signs let successive terms partly cancel, so the partial sums settle to a limit; without the signs ($\sum \frac{1}{n}$) there is no cancellation and the sum grows without bound.