


Solutions

Original authored marking guidance; accept equivalent reasoning and sensible plotting accuracy.

2.1 K s-block, Sc d-block and As p-block. He is s-block because its electrons occupy $1s^2$; the column reflects its full outer shell and chemical behaviour, not a p-electron configuration.

2.2



Linear axes labelled with Z and energy/unit and a consistent scale; all linear points correctly placed. Log axis labelled and scaled, all transformed points correctly placed. He, Ne, Ar and Kr identified as the noble-gas peaks; same ordered data on both plots with no substitution of successive ionisations for different atoms. Accept a correctly labelled logarithmically scaled energy axis.

3.1 $\text{He} \rightarrow \text{Li}$, $\text{Ne} \rightarrow \text{Na}$ and $\text{Ar} \rightarrow \text{K}$. The new outer shell puts the electron further from the nucleus with increased shielding; this outweighs the increase in nuclear charge, so less energy removes it. A log axis compresses the large numerical range and makes proportional changes easier to compare.

3.2 Be $[\text{He}]2s^2$; B $[\text{He}]2s^22p^1$: the B electron is removed from the higher-energy 2p sub-shell, outweighing increased nuclear charge. N $[\text{He}]2s^22p^3$ has singly occupied p orbitals; O $[\text{He}]2s^22p^4$ has a first pair, whose repulsion makes one electron easier to remove. No new occupied shell is added in these comparisons.

3.3 Li/Be metallic; B/C giant covalent; $\text{N}_2/\text{O}_2/\text{F}_2$ simple molecular and Ne monatomic. Carbon melting requires overcoming many strong covalent bonds in its giant structure. Melting nitrogen overcomes attractions between N_2 molecules, retaining the covalent triple bonds inside each molecule. The diagram of a molecule does not imply a giant covalent lattice.

3.4 Valid internal decreases include $\text{Be} \rightarrow \text{B}$, $\text{N} \rightarrow \text{O}$, $\text{Mg} \rightarrow \text{Al}$, $\text{P} \rightarrow \text{S}$, $\text{Ti} \rightarrow \text{V}$, $\text{Fe} \rightarrow \text{Co}$, $\text{Co} \rightarrow \text{Ni}$, $\text{Zn} \rightarrow \text{Ga}$ and $\text{As} \rightarrow \text{Se}$. Melting involves the solid's bonding/structure and attractions be-

tween its particles, rather than removal of one electron from gaseous atoms; first ionisation alone does not determine it.