

# Notes

Pearson Edexcel · International GCSE · Chemistry

## Pearson Edexcel International GCSE · Chemistry: teaching notes

Version: 4CH1 linear qualification; acquired Issue 3 (2024); first examination 2019

This original focus package is partial. It does not certify whole-specification coverage or a reviewed interactive bank.

### Assessment and course boundaries


- This is linear 4CH1, with C-suffixed separate-Chemistry statements retained.
- Paper 1C: 110 marks, 2 h, 61.1%; Paper 2C: 70 marks, 1 h 15 min, 38.9%; no tiers.
- Use specification practical statements, not the UK GCSE core-practical list.

### Particles, bonding and bulk properties

Official-unit focus: 1 Principles of chemistry

A salt crystal conducts when dissolved but not when solid. The ions exist in both states; their ability to move changes.

Ionic bonding is electrostatic attraction between oppositely charged ions. A covalent bond involves shared electrons. Metallic bonding involves attraction between positive metal ions and delocalized electrons.



Alternating charges in an ionic lattice

To explain a bulk property, name the structure, particles, forces and mobile charge carriers. Simple molecular substances can have strong covalent bonds inside molecules but weak attractions between molecules.

Compare substances using evidence such as melting point, conductivity when solid and molten, and solubility. One property rarely proves a structure; use a pattern of evidence.

## Checked worked case

Known: an element has atomic number 12 and mass number 24. Protons = 12; neutrons = mass number - atomic number =  $24 - 12 = 12$ . A 2+ ion has electrons =  $12 - 2 = 10$ . Charge changes electron count, not the nucleus.

## Common error


Melting a simple molecular substance usually overcomes intermolecular attractions; it does not require breaking all covalent bonds within each molecule.

# Amounts, equations and limiting reagents

Official-unit focus: 1 Principles of chemistry

The smallest mass of reactant is not necessarily the limiting reagent. The balanced equation compares particle amounts, not grams directly.

The mole measures amount of substance. Use molar mass to convert mass into amount. Balanced equation coefficients give mole ratios; they do not give equal masses.



Calculate the amount available for each reactant and divide by its coefficient. The smaller ratio limits the reaction. Use that reactant to calculate the maximum product before comparing actual yield.

Write the balanced equation first, include units in molar masses, then convert each given mass or solution volume into amount. Convert cubic centimetres to cubic decimetres before using concentration in moles per cubic decimetre.

## Checked worked case

Known: 2.0 g of Mg reacts with excess acid. Use amount = mass/molar mass. With molar mass Mg = 24.0 g per mole, amount Mg =  $2.0/24.0 = 0.0833$  mol. In  $\text{Mg} + 2\text{HCl} \rightarrow \text{MgCl}_2 + \text{H}_2$ , amount  $\text{H}_2$  = amount Mg = 0.0833 mol.

## Common error


Excess acid means acid does not limit the stated calculation. A coefficient of 2 before HCl does not double the hydrogen amount.

## Redox and electrolysis

Official-unit focus: 1 Principles of chemistry

An aqueous salt solution can produce different electrode products from the molten salt. Water introduces competing species into the system.

Oxidation is loss of electrons and reduction is gain of electrons. In electrolysis, cations move toward the cathode and anions toward the anode. Reduction occurs at the cathode.



Predict products using the specified electrolyte and electrode material. In an aqueous solution, hydrogen or oxygen may form because water-related species compete. Molten salts contain only the ions of the salt.

Use a low-voltage direct-current supply, approved electrodes, and the school risk assessment. Collect gases only by an approved method. Keep chlorine demonstrations

teacher-controlled; do not ask students to generate hazardous gases independently.

## Checked worked case

Known: a copper ion gains two electrons. Half-equation:  $\text{Cu}^{2+} + 2\text{e}^- \rightarrow \text{Cu}$ . One mole of  $\text{Cu}^{2+}$  requires two moles of electrons. For 0.050 mol of copper, electron amount =  $2 \times 0.050 = 0.100$  mol.

## Common error


Electrode signs depend on the cell type. In an electrolytic cell the cathode is negative; reduction remains the defining process at a cathode in every cell.

## Chemical tests and analytical confidence

Official-unit focus: 2 Inorganic chemistry

A coloured flame is useful evidence, but a contaminated wire can give a misleading result. Analytical conclusions depend on the method and controls.

Chemical analysis identifies substances or measures amount. Chromatography separates components because they distribute differently between stationary and mobile phases. A pure substance has characteristic physical properties under stated conditions.



R<sub>f</sub> is distance travelled by a component divided by distance travelled by the solvent front, both measured from the baseline. Compare under the same conditions; an R<sub>f</sub> value alone does not establish identity across different solvents.

Use pencil for the baseline, keep spots above solvent level, mark the solvent front promptly, and run known references alongside unknowns. For ion tests, use clean equipment and separate aliquots to avoid carrying reagents into later tests.

## Checked worked case

Known: spot travels 3.0 cm; solvent front travels 5.0 cm.  $R_f = \text{spot distance} / \text{front distance}$ .  $R_f = 3.0 / 5.0 = 0.60$ .  $R_f$  has no unit. Two matching  $R_f$  values support identification only when other evidence and conditions agree.

## Common error


A single spot can conceal substances that co-elute. Ink on the baseline can dissolve and create extra spots.

## Calorimetry and chemical energy

Official-unit focus: 3 Physical chemistry

A cup warms when two solutions react. The temperature rise measures energy transferred to the surroundings; it does not directly equal the enthalpy change.

Exothermic reactions transfer energy to surroundings. Endothermic reactions take energy from surroundings. Bond breaking requires energy; bond formation releases energy.



Calculate energy transferred from the mass, specific heat capacity and temperature rise when these are given. At GCSE, use reaction profiles and bond-energy differences where your tier specifies them; this lesson does not introduce advanced molar enthalpy calculations.

Use insulation and a lid, measure starting temperatures consistently, stir, and record a temperature-time series. Estimate the reaction temperature from an appropriate extrapolation rather than ignoring cooling during measurement.

## Checked worked case

Known: 100 g solution rises by 5.0 °C; specific heat capacity is 4.18 J per gram per degree.  $q = mc\Delta T$ .  $q = 100 \times 4.18 \times 5.0 = 2,090$  J. This is energy gained by the solution. Insulation and controlled starting temperatures improve the estimate.

## Common error


Heat loss usually lowers the observed temperature rise. The solution gaining heat and the reaction losing heat have opposite signs.

## Rates, catalysts and reliable endpoints

Official-unit focus: 3 Physical chemistry

A faster reaction finishes sooner, but it need not make more product. Rate and final yield answer different questions.

Reaction rate describes reactant used or product formed per time. Higher temperature increases the fraction of collisions with enough energy. A catalyst provides an alternative pathway with lower activation energy.



A product-time graph has a steeper gradient where rate is larger. A tangent estimates instantaneous rate; a secant gives average rate over an interval. The final plateau reflects the total collected product under the stated conditions.

For gas production, check apparatus for leaks, start timing consistently and record volume at regular intervals. Keep concentration, reactant amount and surface area controlled when changing temperature.

## Checked worked case

Known: gas volume increases from 10 to 34 cubic centimetres between 20 and 60 s. Average rate = change in volume/change in time. Rate =  $(34 - 10)/(60 - 20) = 0.60$  cubic centimetres per second. This is not necessarily the instantaneous rate at 40 s.

## Common error


A catalyst does not change the equilibrium constant at a fixed temperature. A mass-loss method cannot detect all reactions, and losing gas through a leak biases a collection experiment.

## Equilibrium and changing conditions

Official-unit focus: 3 Physical chemistry

A reversible reaction can continue in a closed vessel while measured concentrations stay constant. Constant composition does not mean particles have stopped reacting.

Dynamic equilibrium occurs in a closed system when forward and reverse rates are equal. Reactant and product concentrations are constant, but they need not be equal.



Changing conditions can change the equilibrium composition. A catalyst speeds both directions and shortens the time to equilibrium; it does not change the final equilibrium composition at a fixed temperature.

State the balanced equation and whether the forward reaction is exothermic before predicting a temperature effect. Count gas coefficients when considering pressure; pressure has no composition effect when gaseous amounts are equal on both sides.

## Checked worked case

Known: in a closed system, ten forward reaction events and ten reverse events occur in one second. Net change in the product count = forward events - reverse events =  $10 - 10 = 0$ . The measured composition stays constant while reactions continue.

## Common error


Do not use a catalyst to claim a larger equilibrium yield. For heterogeneous equilibria, pure solids are omitted from the usual equilibrium expression.

## Organic structures and reaction pathways

Official-unit focus: 4 Organic chemistry

Two compounds can have the same molecular formula but different structures. Their functional groups help predict which reactions they undergo.

A homologous series shares a functional group and general formula. Structural isomers share a molecular formula but differ in atom connections. Alkenes contain a carbon-carbon double bond.



Distinguish addition, substitution, oxidation and polymerization by tracing bonds before and after reaction. Conditions and reagents belong to the reaction arrow; they are not interchangeable labels.

Draw displayed or structural formulae with the correct number of bonds at each carbon. Use a carbon count to check a proposed synthesis. At advanced level, track reagents and conditions through multistep routes.

## Checked worked case

Known: ethene adds bromine across its double bond. The two-carbon skeleton stays intact and each carbon gains one bromine atom, giving 1,2-dibromoethane. One mole of ethene reacts with one mole of bromine in this addition reaction.

## Common error

Bromine decolourization provides evidence of unsaturation in an appropriate test. It is not proof that an unknown sample is specifically ethene.