

# Reproduction · supported focus

International Baccalaureate · IB Diploma · Biology · SL

## Scope and prerequisites

---

Supported SL focus. First assessment 2025; current subject brief acquired; full Biology guide not acquired. Remaining guide, assessment and practical requirements retain their recorded holds.

Prerequisites: read the stated quantities and units, use arithmetic and the model conditions below. Each lesson develops its own method before independent transfer.

These are original or explicitly fictional teaching examples, not actual measurements or completed assessed learner investigations.

## Sexual reproduction: meiosis and fertilization have different roles


---

### ■ What would explain this observation?

- Gametes normally contain one set of chromosomes, while **fertilization** joins two sets. Maintaining chromosome number across generations requires both processes.
  - Start with a prediction. State the quantities or features you would compare, then decide what evidence could distinguish two explanations.
- 

### ■ Build the model

- Meiosis reduces chromosome number when producing **haploid** cells from a diploid starting cell. Homologous chromosomes separate in meiosis I, while sister chromatids separate in meiosis II. Fertilization combines haploid gamete nuclei to form a diploid zygote. Sexual reproduction can create new combinations of existing alleles.
- haploid: Having one chromosome set in the stated life cycle; fertilization: Fusion of gamete nuclei to form a zygote.



---

### ■ Choose evidence that can test it

- Distinguish homologous chromosomes from sister chromatids, and chromosome number from DNA quantity. Replication before meiosis copies DNA without doubling the number of chromosome sets. Crossing over and independent assortment contribute to genetic variation, while random fertilization further changes combinations. These processes do not make every offspring genetically distinct under every conceivable condition.
- Use a labelled model organism with a stated chromosome number. Track chromosome sets through replication, meiosis I, meiosis II and fertilization. Use anonymous model data rather than personal family or reproductive-health information. Compare organism life cycles carefully; flowering-plant and animal reproductive structures require their own additional teaching.

---

### ■ Work from known quantities

- State the known values and their units. Choose the relation because its assumptions fit this case, then rearrange before substitution.
- Known: a model organism has diploid chromosome number  $2n=12$ . A gamete has  $n=6$ . After fertilization, the zygote has 12 chromosomes. With six independently assorting chromosome pairs and ignoring crossing over, there are  $2^6=64$  possible maternal/paternal chromosome combinations in a gamete. This is a model count, not a probability of an individual phenotype.

#### Example:

A model organism has diploid chromosome number 20. How many chromosomes are normally in a gamete? Use the same sequence: known quantities → model → relation → substitution → unit and interpretation.

## ■ Check the conclusion and its limits

- A gamete is not made by halving the size of an ordinary cell. Independent assortment counts assume distinct homologous alternatives and do not include crossing-over outcomes. Human and plant reproduction cannot be fully replaced by a monohybrid Punnett grid.
- Return to the original observation. Explain what the result supports, which conditions it assumes, and one way to test a competing explanation.

### **Warn:**

DNA replication alone changes a diploid cell into a tetraploid cell. This claim is false: A gamete is not made by halving the size of an ordinary cell. Independent assortment counts assume distinct homologous alternatives and do not include crossing-over outcomes. Human and plant reproduction cannot be fully replaced by a monohybrid Punnett grid.

### **Key:**

Sexual reproduction: meiosis and fertilization have different roles: Distinguish homologous chromosomes from sister chromatids, and chromosome number from DNA quantity. Replication before meiosis copies DNA without doubling the number of chromosome sets. Crossing over and independent assortment contribute to genetic variation, while random fertilization further changes combinations. These processes do not make every offspring genetically distinct under every conceivable condition.