

Calculus · supported focus

International Baccalaureate · IB Diploma · Mathematics:
Applications and Interpretation · SL

Scope and prerequisites

Supported SL focus. First assessment 2021; current through 2028. First-assessment-2029 course is separate.. Remaining guide, assessment and practical requirements retain their recorded holds.

Prerequisites: read the stated quantities and units, use arithmetic and the model conditions below. Each lesson develops its own method before independent transfer.

These are original or explicitly fictional teaching examples, not actual measurements or completed assessed learner investigations.

Derivatives and stationary points

■ What is the slope at one point?

- A curved road has different slopes at different positions. An average gradient cannot describe every point.
- This lesson studies **derivative** : The instantaneous rate of change, also the gradient of a tangent.

■ Choose the mathematical structure

- For $y=ax^n$, $dy/dx=nax^{n-1}$. A stationary point satisfies $dy/dx=0$. Check the sign change of the derivative, or the second derivative when it is nonzero, to classify it.
- State the allowed inputs and units before calculating. An equation should express the relationship, not just record a calculator entry.


■ Work through a checked case

$$\frac{d}{dx}(ax^n) = nax^{n-1}, \quad f'(x) = 0$$

- Check the result against the starting quantities. Substitute into the original relation, or compare the graph and numerical answer where appropriate.

Example:

For $y=x^3-3x$, $dy/dx=3x^2-3$. At $x=1$, the gradient is 0 and $y=-2$. The second derivative is $6x$, positive at $x=1$, so this is a local minimum. At $x=-1$, $y=2$ and the second derivative is negative, giving a local maximum.



■ Test a tempting shortcut

- A zero derivative does not always mean a maximum or minimum: $y=x^3$ is stationary at 0 but continues increasing. An endpoint can also produce an extreme value on a restricted domain.
- When a shortcut fails, identify the assumption it breaks. Keep an exact value until the requested final rounding.

Warn:

Every point with zero derivative is a local maximum or minimum. This claim is false. Explain which definition or assumption it violates.

■ Interpret a new situation

- At GCSE/IGCSE use only the polynomial scope allowed by the tier; do not add chain, product or quotient rules there. At advanced level, connect the derivative to rates and optimization with a valid domain.
- A complete solution gives the mathematical result and explains what it means. Check that it is possible in the stated context.

■ Use this in your course

- Current first-assessment-2021 Applications and Interpretation SL. This is authored concept support; the full guide is needed to certify every objective.
- Give the method before the final answer, and use the paper's calculator and formula rules. Review a wrong answer by locating the first invalid step.

Key:

The instantaneous rate of change, also the gradient of a tangent. Choose the relationship, show the method, check its assumptions and interpret the result.

Root finding and numerical integration

■ How reliable is an approximation?

- A nonlinear equation has no convenient exact solution. An approximation still needs evidence that it is reliable.
- This lesson studies **iteration** : A repeated update in which each new approximation is calculated from the previous one.

■ Choose the mathematical structure

- A continuous function with opposite signs at two endpoints has a root between them. Newton's method uses $x_{\text{next}} = x - f(x)/f'(x)$, with a nonzero derivative. The trapezium rule approximates a definite integral using endpoint heights.
- State the allowed inputs and units before calculating. An equation should express the relationship, not just record a calculator entry.


■ Work through a checked case

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

- Check the result against the starting quantities. Substitute into the original relation, or compare the graph and numerical answer where appropriate.

Example:

For $f(x) = x^2 - 2$ and $x_0 = 1.5$, Newton gives $x_1 = 1.5 - (2.25 - 2)/3 = 1.4166667$. With $y = x^2$ on $[0, 2]$ and two equal strips, $h = 1$ and trapezium area $= (1/2)(0 + 2 \times 1 + 4) = 3$; the exact area is $8/3$.



■ Test a tempting shortcut

- A sign change across a discontinuity does not prove a root. Iteration can diverge or cycle. The trapezium rule's overestimate or underestimate depends on curvature, not just whether the function increases.
- When a shortcut fails, identify the assumption it breaks. Keep an exact value until the requested final rounding.

Warn:

Every sign change proves a root, including one across a discontinuity. This claim is false. Explain which definition or assumption it violates.

■ Interpret a new situation

- Give a stopping criterion and report sensible accuracy. Confirm the approximated root by a sign bracket around the stated rounded answer; explain any failure of the chosen iteration.
- A complete solution gives the mathematical result and explains what it means. Check that it is possible in the stated context.

■ **Use this in your course**

- Current first-assessment-2021 Applications and Interpretation SL. This is authored concept support; the full guide is needed to certify every objective.
- Give the method before the final answer, and use the paper's calculator and formula rules. Review a wrong answer by locating the first invalid step.

Key:

A repeated update in which each new approximation is calculated from the previous one. Choose the relationship, show the method, check its assumptions and interpret the result.